

Table 1.6A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
TOTAL	8,383 ^a	9,334	2,847	3,351	5,535	5,983
18	151	192	30 ^a	74	120	118
19	127	134	46	37	81	97
20	124	166	37	48	87	118
21	112 ^a	184	45	57	67 ^a	127
22	126	176	46	67	80	109
23	169	182	47	51	122	131
24	153	151	56	58	97	93
25	163	148	43	53	120	96
26-29	816	709	274	260	541	449
30-34	1,011	1,150	355	438	656	712
35-39	1,007	897	329	289	677	608
40-44	995	1,082	377	394	618	688
45-49	1,029	1,293	382	526	647	767
50-54	873	1,023	317	268	556	755
55-59	666	913	201	241	465	671
60-64	449	511	118	261	330	249
65 or Older	413	422	143	228	269	194

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.6B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Percentages, 2009 and 2010

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
TOTAL	3.7	4.1	2.6	3.0	4.7	5.1
18	3.0	4.1	1.2 ^a	2.9	5.1	5.4
19	2.8	3.1	1.9	1.7	3.7	4.5
20	3.0	3.7	1.7	2.0	4.4	5.4
21	2.7 ^a	4.3	2.2	2.7	3.2 ^a	6.0
22	3.1	4.3	2.3	3.2	3.9	5.4
23	4.2	4.5	2.4	2.5	6.0	6.6
24	3.7	3.7	2.9	2.9	4.5	4.5
25	4.5	3.8	2.4	2.8	6.5	4.7
26-29	4.9	4.3	3.3	3.1	6.4	5.5
30-34	5.2	5.8	3.6	4.5	6.8	7.0
35-39	5.1	4.6	3.3	3.0	6.9	6.2
40-44	4.8	5.3	3.7	3.9	5.7	6.7
45-49	4.6	5.8	3.5	4.9	5.6	6.7
50-54	3.9	4.7	2.9	2.5	4.9	6.7
55-59	3.6	4.7	2.2	2.6	4.9	6.6
60-64	2.9	3.1	1.6	3.2	4.1	3.0
65 or Older	1.1	1.1	0.9	1.3	1.2	0.9

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.7A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	8,383 ^a	9,334	1,124 ^b	1,334	4,858	5,132	2,401	2,868
GENDER								
Male	2,847	3,351	350 ^a	444	1,718	1,908	780	999
Female	5,535	5,983	774	890	3,140	3,224	1,621	1,869
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	7,486	8,304	965	1,089	4,335	4,573	2,186	2,642
White	6,341	6,743	723 ^a	851	3,632	3,534	1,986	2,359
Black or African American	719 ^a	1,028	147	150	451 ^b	752	121	126
American Indian or Alaska Native	53	87	12	9	37	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	3	*	*	*	*
Asian	164	232	52	25	113	164	*	*
Two or More Races	183	196	24	51	84	76	75	*
Hispanic or Latino	897	1,030	159 ^a	245	523	558	214	226
CURRENT EMPLOYMENT								
Full-Time	3,204	3,499	297 ^a	386	2,247	2,301	660	813
Part-Time	1,123	1,357	327	409	644	700	152	248
Unemployed	804	926	199	198	439	526	166	201
Other ¹	3,251	3,551	301	341	1,527	1,605	1,423	1,606

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.7B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Percentages, 2009 and 2010

Demographic Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	3.7	4.1	3.3 ^a	3.9	4.9	5.2	2.5	3.0
GENDER								
Male	2.6	3.0	2.1	2.6	3.5	3.9	1.8	2.2
Female	4.7	5.1	4.6	5.3	6.3	6.5	3.2	3.6
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	3.8	4.2	3.5	4.0	5.3	5.6	2.5	3.0
White	4.1	4.3	3.6 ^a	4.2	5.8	5.7	2.8	3.2
Black or African American	2.7 ^a	3.9	3.1	3.1	3.7 ^b	6.1	1.3	1.3
American Indian or Alaska Native	4.8	8.1	5.3	4.8	8.3	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	2.1	*	*	*	*
Asian	1.6	2.2	3.4	1.7	2.1	3.1	*	*
Two or More Races	7.5	7.4	5.5	9.4	7.9	7.3	8.0	*
Hispanic or Latino	2.9	3.2	2.6 ^a	3.7	3.1	3.3	2.7	2.7
CURRENT EMPLOYMENT								
Full-Time	2.8	3.1	2.4 ^a	3.1	3.4	3.5	1.8	2.2
Part-Time	3.5	4.1	3.5	4.3	5.4	5.7	1.4	2.2
Unemployed	5.4	6.3	4.4	4.4	6.2	7.9	5.2	5.6
Other ¹	4.9	5.3	4.1	4.5	10.9	11.4	3.2	3.5

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.8A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic/Socioeconomic Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	8,383 ^a	9,334	1,124 ^b	1,334	4,858	5,132	2,401	2,868
GEOGRAPHIC REGION								
Northeast	1,476	1,651	219	258	869	950	389	442
Midwest	2,151	2,276	304	273	1,295	1,157	552 ^a	846
South	2,827	3,113	304 ^b	455	1,656	1,713	866	945
West	1,928	2,293	298	347	1,038	1,312	593	634
COUNTY TYPE								
Large Metro	4,237	4,700	594 ^a	720	2,310	2,670	1,332	1,310
Small Metro	2,551	2,972	345 ^b	455	1,598	1,681	608	836
250K – 1 Mil. Pop.	1,663 ^a	2,091	212 ^b	314	1,075	1,192	377	585
< 250K Pop.	888	880	133	141	523	489	231	251
Nonmetro	1,594	1,662	185	160	949	781	460	721
Urbanized	664	758	81 ^a	52	393	369	190	338
Less Urbanized	741	700	89	92	472	359	180	250
Completely Rural	189	203	16	17	84	53	90	*
POVERTY LEVEL¹								
Less Than 100%	1,792	2,075	275	329	995	1,132	521	614
100-199%	1,993	2,242	287 ^a	369	1,263	1,232	443	641
200% or More	4,551	4,987	514	607	2,600	2,767	1,437	1,613
GOVERNMENT ASSISTANCE²								
Yes	2,571 ^a	3,093	323 ^a	420	1,524	1,754	725	920
No	5,812	6,240	802	914	3,334	3,378	1,676	1,948
HEALTH INSURANCE³								
Private	4,099	4,765	523 ^b	673	2,423	2,545	1,154	1,547
Medicaid/CHIP ⁴	1,680	1,694	255	208	1,012	933	414	553
Other ⁵	1,799	2,204	86	126	660	889	1,053	1,188
No Coverage	1,710	1,909	325	368	1,109	1,210	276	330

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

³ Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

⁴ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁵ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.8B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: Percentages, 2009 and 2010

Geographic/Socioeconomic Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	3.7	4.1	3.3 ^a	3.9	4.9	5.2	2.5	3.0
GEOGRAPHIC REGION								
Northeast	3.5	3.9	3.6	4.3	4.9	5.3	2.2	2.4
Midwest	4.3	4.6	4.1	3.6	6.1	5.5	2.6	3.9
South	3.4	3.7	2.5 ^b	3.7	4.5	4.7	2.5	2.7
West	3.7	4.3	3.7	4.3	4.4	5.6	2.9	3.0
COUNTY TYPE								
Large Metro	3.5	3.8	3.3	4.0	4.2	4.9	2.8	2.6
Small Metro	3.7	4.2	3.3 ^a	4.1	5.5	5.7	2.1	2.9
250K – 1 Mil. Pop.	3.7 ^a	4.5	3.2 ^b	4.5	5.5	5.9	2.0	3.0
< 250K Pop.	3.8	3.7	3.5	3.4	5.4	5.1	2.3	2.6
Nonmetro	4.3	4.5	3.5	3.3	6.4	5.5	2.6	4.1
Urbanized	4.1	4.9	3.2	2.4	6.2	6.1	2.6	4.5
Less Urbanized	4.3	4.0	3.7	3.9	6.7	5.2	2.3	3.0
Completely Rural	5.0	6.0	4.8	4.8	6.7	4.5	4.0	*
POVERTY LEVEL¹								
Less Than 100%	6.5	7.0	3.5	3.8	7.9	9.0	7.3	7.3
100-199%	4.5	4.7	3.6	4.4	6.8	5.9	2.6	3.5
200% or More	2.9	3.3	3.1 ^a	3.8	3.8	4.2	2.1	2.3
GOVERNMENT ASSISTANCE²								
Yes	7.4	7.7	4.9	5.5	9.3	9.4	6.1	6.6
No	3.0	3.3	3.0 ^a	3.5	4.0	4.2	2.0	2.4
HEALTH INSURANCE³								
Private	2.7 ^a	3.2	2.9 ^b	3.7	3.6	3.9	1.7	2.3
Medicaid/CHIP ⁴	8.7	8.0	5.4	4.1	11.7	10.3	6.9	7.9
Other ⁵	3.4	4.1	4.1	5.5	10.3	13.3	2.4	2.6
No Coverage	4.7	4.9	3.4	3.9	5.6	5.7	4.0	4.1

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

³ Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

⁴ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁵ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.